

Longitudinal and transverse structure functions of proton and deuteron at large x^a

G. Ricco^(a) and S. Simula^(b)

^(a)Physics Department, University of Genova and INFN, Sezione di Genova,
Via Dodecanneso 33, I-16146 Genova, Italy

^(b)Istituto Nazionale di Fisica Nucleare, Sezione Roma III,
Via della Vasca Navale 84, I-00146 Roma, Italy

Abstract

Higher-twist effects in the low-order moments of the longitudinal and transverse structure functions of proton and deuteron have been analyzed using available phenomenological fits of existing data in the Q^2 range between 1 and 20 $(GeV/c)^2$. Both twist-4 and twist-6 contributions have been determined adopting the Nachtmann definition of moments, which allows to disentangle properly target-mass effects. The extraction of the matrix elements of the relevant twist-4 operators, describing quark-quark and quark-gluon correlations, is carried out in case of the second moment. The need of transverse data with better quality for $x \gtrsim 0.5$ and $Q^2 \lesssim 10 (GeV/c)^2$ as well as more precise and systematic determinations of the L/T separation make *JLab* @ 12 *GeV* a good place to improve our understanding of the non-perturbative structure of hadrons.

^aTo appear in the Proceedings of the International Workshop on *JLab: Physics and Instrumentation with 6-12 GeV Beams*, Jefferson Lab (Newport News, USA), June 15-18, 1998.

1 INTRODUCTION

The experimental investigation of deep-inelastic lepton-nucleon scattering has provided a wealth of information on the occurrence of Bjorken scaling and its violations, giving a decisive support to the rise of the parton model and its *QCD*-improved version, which properly describe the logarithmic violations to scaling in the asymptotic region. However, in the pre-asymptotic region the full dependence of the nucleon response on the squared four-momentum transfer, Q^2 , is affected also by power-type corrections, which originate from non-perturbative physics and can be analyzed in the framework provided by the Operator Product Expansion (*OPE*). The logarithmic scale dependence is therefore related to the so-called leading twist operators, which in the parton language are one-body operators whose matrix elements yield the contribution of the individual partons to the nucleon response. On the contrary, power-type corrections are related to higher-twist operators measuring the relevance of correlations among partons [1].

In case of unpolarized inelastic electron scattering the nucleon response is described by two independent quantities: the transverse $F_2(x, Q^2)$ and the longitudinal $F_L(x, Q^2)$ structure functions, the latter being related to the ratio of the longitudinal to transverse cross sections, $R_{L/T}(x, Q^2)$, by $F_L(x, Q^2) = F_2(x, Q^2) (1 + 4M^2x^2/Q^2)R_{L/T}(x, Q^2)/[1 + R_{L/T}(x, Q^2)]$, where $x \equiv Q^2/2M\nu$ is the Bjorken variable, M is the nucleon mass and ν is the energy transfer in the nucleon rest frame. Systematic measurements [2] of the transverse function $F_2(x, Q^2)$ for proton and deuteron targets have been carried out in the kinematical range $10^{-4} \lesssim x \lesssim 1$ and for Q^2 values up to several hundreds of $(GeV/c)^2$, while data on the ratio $R_{L/T}(x, Q^2)$ are available for $0.002 \lesssim x \lesssim 0.8$ and $0.5 \lesssim Q^2(GeV/c)^2 \lesssim 70$, though they are still fluctuating and affected by large errors. Consequently, phenomenological fits for both $F_2(x, Q^2)$ and $F_L(x, Q^2)$ are available, but for the latter quantity the interpolation formulae greatly suffer for very weak constraints. The analysis of the world data set [2] has allowed to extract the parton densities in the nucleon, including their *QCD*-predicted logarithmic Q^2 evolution, as well as to signal the presence of power-type scaling violations at *large* x ($\gtrsim 0.7$) and *low* Q^2 ($\lesssim 10 (GeV/c)^2$). The analysis of these kinematical regions, where higher-twist effects are important, represents the aim of the present contribution. A more detailed version of our work will be available soon in [4].

2 TWIST ANALYSIS

An important and effective tool for the theoretical investigation of the complete Q^2 dependence of hadron structure functions is the *OPE*, which leads to the well-known twist expansion for the moments of the structure functions. In our analysis we do not use the Cornwall-Norton definition of the moments, since target-mass corrections, i.e. terms containing powers of M^2/Q^2 , would contribute. Instead of that we will adopt the Nachtmann

definition [3]:

$$M_n^{(T)}(Q^2) \equiv \int_0^1 dx \frac{\xi^{n+1}}{x^3} F_2(x, Q^2) \frac{3 + 3(n+1)r + n(n+2)r^2}{(n+2)(n+3)} \quad (1)$$

$$M_n^{(L)}(Q^2) \equiv \int_0^1 dx \frac{\xi^{n+1}}{x^3} \left[F_L(x, Q^2) + \frac{4M^2 x}{Q^2} F_2(x, Q^2) \frac{(n+1)\xi - 2(n+2)x}{(n+2)(n+3)} \right] \quad (2)$$

where $n \geq 2$, $r \equiv \sqrt{1 + 4M^2 x^2 / Q^2}$ and $\xi \equiv 2x / (1 + r)$ is the Nachtmann variable. Using the *experimental* $F_2(x, Q^2)$ and $F_L(x, Q^2)$ in Eqs. (1-2), target-mass effects are canceled out and therefore the twist expansions of the experimental $M_n^{(T)}(Q^2)$ and $M_n^{(L)}(Q^2)$ contain only *dynamical* twists, namely

$$M_n^{L(T)}(Q^2) = \sum_{\tau=2}^{\infty} C_{n,\tau}^{L(T)}(Q^2/\mu^2) A_{n,\tau}^{L(T)}(\mu^2) (\mu^2/Q^2)^{\frac{\tau-2}{2}} \quad (3)$$

where μ is the renormalization scale, $C_{n,\tau}^{L(T)}(Q^2/\mu^2)$ is a Wilson coefficient calculable within perturbative *QCD* and $A_{n,\tau}^{L(T)}(\mu^2)$ corresponds to the matrix elements of operators of twist τ and spin n . In our analysis the expansion (3) is simplified into

$$\begin{aligned} M_n^{L(T)}(Q^2) = & A_n^{L(T)}(Q^2) + a_n^{(4)}[L(T)] \frac{\mu^2}{Q^2} \left(\frac{\alpha_s(\mu^2)}{\alpha_s(Q^2)} \right)^{\gamma_n^{(4)}[L(T)]} \\ & + a_n^{(6)}[L(T)] \frac{\mu^4}{Q^4} \left(\frac{\alpha_s(\mu^2)}{\alpha_s(Q^2)} \right)^{\gamma_n^{(6)}[L(T)]} \end{aligned} \quad (4)$$

where $\alpha_s(Q^2)$ is the running coupling constant and $A_n^{L(T)}(Q^2)$ is the leading-twist contribution, whose Q^2 dependence is calculated according to the *pQCD* predictions at *NLO*. In Eq. (4) the last two terms in the r.h.s. are simplified parametrizations of the twist-4 and twist-6 contributions, respectively, as suggested in Ref. [5]; the quantities $a_4^n[L(T)]$ ($a_6^n[L(T)]$) and $\gamma_4^n[L(T)]$ ($\gamma_6^n[L(T)]$) represent the effective magnitude and anomalous dimension of twist-4 (twist-6) operators.

3 MAIN RESULTS

Equation (4) has been applied to the fit of the Q^2 behavior of the *experimental* moments $M_n^{L(T)}(Q^2)$ in the range $1 \lesssim Q^2 (\text{GeV}/c)^2 \lesssim 20$. In order to evaluate the r.h.s. of Eqs. (1-2) available phenomenological fits based on the data of Ref. [2] have been used and the elastic peak contributions have been added, as it is required by the inclusive nature of the *OPE*. In case of the deuteron, the nucleon elastic peak leads to the quasi-elastic contribution to the inclusive cross section, which has been evaluated through a convolution approach checked

against available *SLAC* data (see [4] for details). In case of the analysis of the transverse moments, besides the free parameters $a_n^{(\tau)}$ and $\gamma_n^{(\tau)}$ appearing in the twist-4 and twist-6 terms of Eq. (4), also the magnitude of the leading term is simultaneously determined from the fit of our pseudo-data. On the contrary, the parton densities of Ref. [6] have been adopted for evaluating $A_n^L(Q^2)$ (see again [4] for details). Thanks to the decoupling of the singlet-quark and gluon operators at large x , the non-singlet evolution of the leading twist can be safely applied for the analysis of the moments with $n \geq 4$. In our work we have adopted a renormalization scale $\mu = 1 \text{ GeV}$ as in [5].

The main results of our analysis of the transverse moments for both proton and deuteron targets can be summarized as follows: i) the simplified twist expansion (4), containing up to the twist-6 term, is able to reproduce the Q^2 dependence of the transverse moments starting from $Q^2 \simeq 1 \text{ (GeV/c)}^2$; ii) the second moment $M_2^T(Q^2)$ is only slightly affected by the twist-4 and almost unaffected by the twist-6 (see Fig. 1(a)); iii) on the contrary, both twist-4 and twist-6 significantly contribute to the moments of order $n \geq 4$ (see Fig. 1(b)), in accord with the presence of higher-twist effects at large x ; iv) the signs of the twist-4 and twist-6 contributions turn out to be opposite.




Figure 1. Second (a) and fourth (b) moments of the proton transverse structure function versus Q^2 . Open dots: pseudo-data calculated by Eq. (1) using available phenomenological fits of existing data [2] on $F_2(x, Q^2)$; the error bars are obtained using the quoted uncertainties of the phenomenological fits. Solid lines: result of our twist analysis based on Eq. (4); dashed-dotted, dashed and dotted lines: twist-2, twist-4 and twist-6 contributions, respectively.

Basing on naive counting arguments, one can argue that the twist expansion (4) of the transverse moments for $Q^2 \sim \mu^2$ can be rewritten as: $M_n^T(\mu^2) = A_n^T(\mu^2)[1 + n(\mu_n^{(4)}/\mu)^2 - n^2(\mu_n^{(6)}/\mu)^4]$, with $\mu_n^{(\tau)}$ approximately independent of n for $n \gtrsim 4$. Thus, one gets

$$\mu_n^{(4)} = \mu \sqrt{\frac{a_n^{(4)}[T]}{nA_n^T(\mu^2)}}, \quad \mu_n^{(6)} = \mu \left[\frac{|a_n^{(6)}[T]|}{n^2 A_n^T(\mu^2)} \right]^{1/4}. \quad (5)$$

Our results for $\mu_n^{(\tau)}$ are collected in Fig. 2, where it can clearly be seen that the mass scales of the twist-4 and twist-6 terms of our analysis are $\mu_n^{(4)} \simeq \mu^{(4)} \simeq 1 \text{ GeV}$ and $\mu_n^{(6)} \simeq \mu^{(6)} \simeq$

0.6 GeV. The value obtained for $\mu^{(4)}$ is significantly higher than the naive expectation $\mu^{(4)} \simeq \sqrt{\langle k_{\perp}^2 \rangle} \simeq 0.3$ GeV [7] as well as higher than the result of other twist-4 analyses (see [5]).




Figure 2. The mass scale $\mu_n^{(\tau)}$ (Eq. (5)) of the twist-4 and twist-6 terms of our twist analysis (4). Open and full markers correspond to the proton and deuteron case, while dots and squares are our results for the twist-4 and twist-6, respectively.

In case of the longitudinal channel the uncertainties in the calculation of the moments are remarkably larger than those of the transverse ones. The effects of the higher-twists are still dominant in the second moment $M_2^L(Q^2)$ up to Q^2 of several $(\text{GeV}/c)^2$ (see Fig. 3(a)). Note that the moments with $n \geq 4$ can be reproduced by considering the leading twist plus a twist-4 term only (see Fig. 3(b)).




Figure 3. The same as in Fig. 1, but in case of the proton longitudinal structure function.

The main goal of the investigation of higher-twist effects is to disentangle the separate contributions of the various operators of a given twist yielding the relevant multiparton correlations. This is not an easy task, due to the contributions of many operators for any given twist (see [1, 8]). Following Ref. [1], there are seven twist-4 operators contributing to the second moment: three in the non-singlet channel and four in the singlet one; the explicit expressions of these operators can be read off from [1]. In the non-singlet case it is

possible to write down three independent equations using, besides $M_2^T(Q^2)$ and $M_2^L(Q^2)$, the second moment of the structure function $F_3(x, Q^2)$, which can be determined in neutrino and antineutrino scattering experiments, like the recent measurement performed by the *CCFR* collaboration [9] at FermiLab. The neutrino data have been analyzed at *NNLO* in Ref. [10], obtaining a determination of the twist-4 contribution. Using all these experimental results and adopting the notation of Ref. [1], we have got: $A^{NS} = -9.0 \pm 4.5$ and $B^{NS} = -2.0 \pm 0.4$ for the quark-gluon correlation operators, and $C^{NS} = 5.2 \pm 2.8$ for the quark-quark correlation operator. In the singlet case the neutrino data on $F_3(x, Q^2)$ cannot help in determining the various twist-4 operators; assuming that the quark-quark correlation matrix element C^S is the same in neutrino and electron experiments, we have got the following constraints: $A^S + 6B^S = 12.5 \pm 1.8$ and $8C^S + 5A^S - 2B^S = 0$.

Finally, we point out that transverse data with better quality are still needed for $x \gtrsim 0.5$ and $Q^2 \lesssim 10 (GeV/c)^2$ as well as more precise and systematic determinations of the L/T separation are required; for these reasons an electron facility, like *JLab* @ 12 GeV, is a good place to investigate multiparton correlations in the hadron structure.

References

- [1] E.V. Shuryak and A.I. Vainshtein: Nucl. Phys. **B199** (1982) 451; *ibidem* **B201** (1982) 141.
- [2] A. Bodek et al.: Phys. Rev. **D20** (1979) 7. S. Stein et al.: Phys. Rev. **D12** (1975) 1884. J.J. Aubert et al.: Nucl. Phys. **B259** (1985) 189; *ibidem* **B293** (1987) 740. A.C. Benvenuti et al.: Phys. Lett. **B223** (1989) 485; *ibidem* **B237** (1990) 592. M. Arneodo et al.: Nucl. Phys. **B483** (1997) 3197.
- [3] O. Natchmann: Nucl. Phys. **B63** (1973) 237.
- [4] G. Ricco and S. Simula: preprint INFN-RM3 98/2, September 1998.
- [5] X. Ji and P. Unrau: Phys. Rev. **D52** (1995) 72.
- [6] M. Glück, E. Reya and A. Vogt: Z. Phys. **C67** (1995) 433.
- [7] A. De Rujula, H. Georgi and H.D. Politzer: Ann. Phys. **103** (1977) 315. R.K. Ellis, W. Furmanski and R. Petronzio: Nucl. Phys. **B212** (1983) 29.
- [8] R.L. Jaffe and M. Soldate: Phys. Rev. **D26** (1982) 49.
- [9] *CCFR* Collaboration, W.G. Seligman et al.: hep-ex/9701017.
- [10] A.V. Sidorov and M.V. Tokarev: N. Cim. **A110** (1997) 1401.